Gluteus Medius/Minimus Rehabilitation Protocol

Patient may progress to Phase 2 when they have achieved the following: minimal pain with phase 1 exercises, 90 degrees of pain free flexion, minimal range of motion limitations with internal rotation/extension/abduction.

PHASE 2: INTERMEDIATE PHASE Weeks 4-6 Continue with previous or modified versions of previous exercises, but may add: ☐ Crunches ☐ Double leg bridges on the ball ☐ Gradually increase resistance with stationary bike ☐ Standing adduction with theraband resistance Aquatic exercises- flutter kick, swimming, 4 way hip with water weights as tolerated, step ups ☐ Superman in quadruped ☐ Single leg bridges as tolerated Patient may progress to Phase 3 when they have achieved the following: 105 degrees of flexion, 20 degrees of ER, hip flexion strength >60% uninvolved side, adduction/IR/extension/ER strength 70% uninvolved side, pain free with phase 2 exercises PHASE 3: ADVANCED PHASE Continue with previous or modified versions of previous exercises, but may add: ☐ Log rolling ☐ Calf Raises ☐ Physioball exercises – hip lift, bent knee hip lift, hamstring curls, balance Week 8 Continue with previous or modified versions of previous exercises, but may add: ☐ Mini squats ☐ Leg press (start with minimal resistance and increase by patient tolerance) ☐ Step-Up \square Side stepping over cones ☐ Core strengthening on physioball ☐ Abduction isometrics- minimal resistance without pain Week 9 Continue with previous or modified versions of previous exercises, but may add: ☐ Standing abduction without resistance ☐ Elliptical ☐ Seated IR/ER □ Clamshells Week 10 Continue with previous or modified versions of previous exercises, but may add: ☐ Sidelying abduction without resistance as tolerated (instead of standing) ☐ Single leg balance ☐ Abduction and adduction leg raise ☐ BOSU squats

Patient may progress to Phase 4 when they have achieved the following: gluteus medius strength 60-70%, patient can perform phase 3 exercises without pain, pain-free, normal gait pattern.

Weeks 11-15 Continue with previous or modified versions of previous exercises, but may add: ☐ Standing abduction with Theraband resistance as tolerated without pain ☐ Pool running ☐ Lateral step ups ☐ Elliptical ☐ Step drills, quick feet step ups, forward, lateral, carioca ☐ Plyometrics – double leg and single leg shuttle jumps ☐ Theraband walking patterns 1 rep of six exercises at 50 yards – forward, sidestepping, carioca, monster steps, backward, half circles. ☐ Sidestepping with resistance – can use sports cord ☐ Single leg body squats ☐ Lunges – from single plane to tri-planar; add medicine balls for resistance ☐ Sport specific training Patient may progress to final phase when they have achieved the following: single leg mini squat with a level pelvis, can perform phase 4 exercises with proper body mechanics and without pain. **FINAL PHASE** Weeks 16+ Continue with previous or modified versions of previous exercises, but may add: ☐ Running progression ☐ Sport specific drills ☐ Traditional weight training ☐ Plyometric training Criteria for full return to sport: ☐ Full range of motion ☐ Hip strength equal to uninvolved side; single leg pick-up with level pelvis ☐ Ability to perform sport-specific drills at full speed without pain

☐ Restore full gluteus medius strength before higher level activities are added

☐ Completion of functional sports test

PHASE 4: SPORTS SPECIFIC REHAB CLINIC BASED PROGRESSION